


# Summary

Name	IL-6 R alpha/IL-6RA/CD126/IL-6 Receptor Subunit alpha
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Interleukin-6 Receptor Subunit Alpha is produced by our Mammalian expression system and the target gene encoding Leu20-Asp358 is expressed with a 6His tag at the C-terminus.
Accession #	P08887
Host	Human Cells
Species	Human
Predicted Molecular Mass	38.9 KDa
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

# **SDS-PAGE** image



## Background



# Alternative NamesInterleukin-6 receptor subunit alpha; IL-6R subunit alpha; IL-6R-alpha; IL-6R 1;<br/>Membrane glycoprotein 80; gp80; CD126BackgroundInterleukin 6 is a potent pleiotropic cytokine that regulates cell growth and<br/>differentiation and plays an important role in the immune response. IL6Ra is a part<br/>of the receptor for interleukin 6 cytokine. IL6Ra binds to IL6 with low affinity, but<br/>does not transduce a signal. Signal activation necessitates an association with<br/>IL6ST. Activation may lead to the regulation of the immune response, acute-phase<br/>reactions and hematopoiesis. Low concentration of a soluble form of IL6 receptor<br/>acts as an agonist of IL6 activity. Dysregulated production of IL6 and this receptor<br/>are implicated in the pathogenesis of many diseases, such as multiple myeloma,<br/>autoimmune diseases and prostate cancer.

### Note

For Research Use Only, Not for Diagnostic Use.