


## Summary

Name	IL-9/Interleukin-9
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Interleukin-9 is produced by our Mammalian expression system and the target gene encoding GIn19-IIe144 is expressed with a 6His tag at the C-terminus.
Accession #	P15248
Host	Human Cells
Species	Human
Predicted Molecular Mass	15.16 KDa
Formulation	Lyophilized from a 0.2 $\mu m$ filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## **SDS-PAGE** image



## Background



Alternative Names	Interleukin-9; IL-9; Cytokine P40; T-Cell Growth Factor P40; IL9
Background	Interleukin-9 (IL-9) is a secreted protein that belongs to the IL-7/IL-9 family. IL-9 supports IL-2 independent and IL-4 independent growth of helper T-cells. IL-9 stimulates cell proliferation and prevents apoptosis. It functions through the IL-9 receptor (IL-9R), which activates different signal transducer and activator (STAT) proteins and thus connects this cytokine to various biological processes. IL-9 has been identified as a candidate gene for asthma. IL-9 is a determining factor in the pathogenesis of bronchial hyperresponsiveness.

## Note

For Research Use Only, Not for Diagnostic Use.