Product Name: Recombinant Human ITGA5 (C-6His)

Catalog #: PHH0960



Summary

Name Integrin alpha-5/CD49e

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Human Integrin α -5 is produced by our Mammalian expression

system and the target gene encoding Phe42-Tyr995 is expressed with a 6His

tag at the C-terminus.

Accession # P08648

Host **Human Cells**

Species Human

Predicted Molecular Mass 105.11 KDa

Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2. **Formulation**

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 Stability&Storage


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human ITGA5 (C-6His)

Catalog #: PHH0960



Background

Alternative Names

Integrin Alpha-5; CD49 Antigen-Like Family Member E; Fibronectin Receptor Subunit Alpha; Integrin Alpha-F; VLA-5; CD49e; ITGA5; FNRA

Background

Integrin $\alpha\text{-}5$ belongs to the Integrin α chain family and contains 7 FG-GAP repeats. Integrin $\alpha\text{-}5$ joins with Integrin- β1 to form a fibronectin and laminin receptor which recognizes the sequence R-G-D in its ligands. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposis sarcoma lesions. It is expressed on fibroblasts, endothelial cells, peripheral T cells and platelets. Integrin $\alpha\text{-}5$ undergoes post-translational cleavage in the extracellular domain to yield disulfide-linked light and heavy chains. In addition to adhesion, ITGA5 participates in cell-surface mediated signalling.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838