


Summary

Name	LRRTM2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Leucine-Rich Repeat Transmembrane Neuronal Protein 2 is produced by our Mammalian expression system and the target gene encoding Cys34-Arg422 is expressed with a 6His tag at the C-terminus.
Accession #	O43300
Host	Human Cells
Species	Human
Predicted Molecular Mass	45.55 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human LRRTM2 (C-6His)
Catalog #: PHH1102



Alternative Names

Leucine-Rich Repeat Transmembrane Neuronal Protein 2; Leucine-Rich Repeat Neuronal 2 Protein; LRRTM2; KIAA0416; LRRN2

Background

Leucine-Rich Repeat Transmembrane Neuronal Protein 2 (LRRTM2) is a single-pass type I membrane protein that belongs to the LRRTM family. It contains ten LRR (leucine-rich) repeats, one LRRCT domain, and one LRRNT domain. LRRTM2 is expressed in neuronal tissues, and it interacts with DLG4 and NRXN1. LRRTM2 has been suggested to be involved in the development and maintenance of excitatory synapses in the vertebrate nervous system. LRRTM2 also regulates the surface expression of AMPA receptors. LRRTM2 acts as a ligand for the presynaptic receptors NRXN1-A and NRXN1-B.

Note

For Research Use Only , Not for Diagnostic Use.