


Summary

Name	CD58/LFA-3/Lymphocyte Function-associated Antigen 3
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Lymphocyte Function-associated Antigen 3 is produced by our Mammalian expression system and the target gene encoding Phe29-Arg215 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	AAH05930
Host	Human Cells
Species	Human
Predicted Molecular Mass	48.6 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human CD58 (C-Fc)
Catalog #: PHH1115



Alternative Names

Lymphocyte Function-Associated Antigen 3; Surface Glycoprotein LFA-3; CD58; LFA3; Ag3; CD58 antigen

Background

Lymphocyte function-associated antigen 3 (LFA-3/CD58) is a single-pass type I membrane protein. CD58 is widely expressed on hematopoietic and non-hematopoietic human tissue and has been found on leukocytes, erythrocytes, endothelial cells, epithelial cells and fibroblasts of human origin. It is a Ligand of the T-lymphocyte CD2 glycoprotein. This interaction is important in mediating thymocyte interactions with thymic epithelial cells, antigen-independent and -dependent interactions of T-lymphocytes with target cells and antigen-presenting cells and the T-lymphocyte rosetting with erythrocytes. In addition, the LFA-3/CD2 interaction may prime response by both the CD2+ and LFA-3+ cells.

Note

For Research Use Only , Not for Diagnostic Use.