


## Summary

| Name                     | MFAP4/Microfibril-associated glycoprotein 4  |
|--------------------------|--|
| Purity                   | Greater than 95% as determined by reducing SDS-PAGE  |
| Endotoxin level          | <1 EU/µg as determined by LAL test.  |
| Construction             | Recombinant Human Microfibril-associated Glycoprotein 4 is produced by our Mammalian expression system and the target gene encoding Val22-<br>Ala255 is expressed with a 6His tag at the C-terminus.   |
| Accession #              | AAH62415.1   |
| Host                     | Human Cells  |
| Species                  | Human  |
| Predicted Molecular Mass | 27.5 KDa   |
| Formulation              | Lyophilized from a 0.2 $\mu m$ filtered solution of 20mM PB, 150mM NaCl, pH 7.4.   |
| Shipping                 | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.   |
| Stability&Storage        | Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.   |
| Reconstitution           | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is<br>not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve<br>the lyophilized protein in distilled water. Please aliquot the reconstituted solution<br>to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not<br>mix by vortex or pipetting. It is not recommended to reconstitute to a<br>concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled<br>water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

## **SDS-PAGE** image



## Background



Alternative NamesMicrofibril-associated glycoprotein 4BackgroundMicrofibril-associated glycoprotein 4(MFAP4) is a secreted protein and contains 1<br/>fibrinogen C-terminal domain, similarity to a bovine microfibril-associated protein.<br/>The protein has binding specificities for both collagen and carbohydrate. It is<br/>thought to be an extracellular matrix protein which is involved in calcium-<br/>dependent cell adhesion or intercellular interactions. The gene is located within the<br/>Smith-Magenis syndrome region.

## Note

For Research Use Only, Not for Diagnostic Use.