


Summary

Name	NCR1/NKp46/CD335/Natural Cytotoxicity Triggering Receptor 1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Natural Cytotoxicity Triggering Receptor 1 is produced by our Mammalian expression system and the target gene encoding Gln22-Asn254 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	AAH64806
Host	Human Cells
Species	Human
Predicted Molecular Mass	53.5 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human NCR1 (C-Fc)
Catalog #: PHH1192



Background

Alternative Names

Natural cytotoxicity triggering receptor 1; Lymphocyte antigen 94 homolog; NK cell-activating receptor; Natural killer cell p46-related protein; NK-p46; NKp46; hNKp46; CD335; NCR1; LY94

Background

Natural cytotoxicity triggering receptor 1(NCR1) is a single-pass type I membrane protein .It contains 2 Ig-like (immunoglobulin-like) domains and belongs to the natural cytotoxicity receptor (NCR) family. The protein is a natural killer (NK) lymphocyte-activating receptor. It is involved in major aspects of NK immune function and shows a high degree of lineage specificity in blood and bone marrow.

Note

For Research Use Only , Not for Diagnostic Use.