


Summary

Name	NKG2D Ligand 2/NKG2DL2/N2DL2/ULBP-2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human UL16 Binding Protein-2/NKG2D Ligand 2 is produced by our Mammalian expression system and the target gene encoding Gly26-Ser217 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q9BZM5
Host	Human Cells
Species	Human
Predicted Molecular Mass	48.87 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human NKG2DL2(C-Fc)
Catalog #: PHH1232



Alternative Names

NKG2D Ligand 2; N2DL-2; NKG2DL2; ALCAN-Alpha; Retinoic Acid Early Transcript 1H; UL16-Binding Protein 2; ULBP2; N2DL2; RAET1H

Background

NKG2D Ligand 2 (N2DL2) is a member of a family of cell-surface proteins. N2DL2 function as ligands for human cytomegalovirus glycoprotein UL16. N2DL2 is anchored to the membrane via a GPI-linkage. N2DL2 is bind to human NKG2D, an activating receptor expressed on NK cells, NKT cells, T cells. Engagement of NKG2D results in the activation of cytolytic activity and cytokine production by these effects cells. The ULBPs are expressed on some tumor cells and have been implicated in tumor surveillance.

Note

For Research Use Only , Not for Diagnostic Use.