


Summary

Name	NTAL/LAT2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Linker for Activation of T-Cells Family Member 2 is produced by our Mammalian expression system and the target gene encoding Arg27-Ala243 is expressed with a 6His tag at the C-terminus.
Accession #	Q9GZY6
Host	Human Cells
Species	Human
Predicted Molecular Mass	25 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human NTAL (C-6His)
Catalog #: PHH1243



Alternative Names

Linker for Activation of T-Cells Family Member 2; Linker for Activation of B-Cells; Membrane-Associated Adapter Molecule; Non-T-Cell Activation Linker; Williams-Beuren Syndrome Chromosomal Region 15 Protein; Williams-Beuren Syndrome Chromosomal Region 5 Protein; LAT2; LAB; NTAL; WBS15; WBSCR15; WBSCR5

Background

Linker for Activation of T-Cells Family Member 2 (LAT2) is a single-pass type III membrane protein. LAT2 is highly expressed in the spleen, peripheral blood lymphocytes, and germinal centers of lymph nodes. LAT2 is involved in FCER1 (high affinity immunoglobulin epsilon receptor)-mediated signaling in mast cells. It may also be involved in BCR (B-cell antigen receptor)-mediated signaling in B-cells and FCGR1 (high affinity immunoglobulin gamma Fc receptor I)-mediated signaling in myeloid cells. Coupling activate of these receptors and their associated kinases with distal intracellular events through the recruitment of GRB2.

Note

For Research Use Only , Not for Diagnostic Use.