


Summary

Name	PRKG1/cGMP-dependent protein kinase 1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human cGMP-Dependent Protein Kinase 1 is produced by our Mammalian expression system and the target gene encoding Gly2-Phe686 is expressed with a 6His tag at the C-terminus.
Accession #	Q13976-2
Host	Human Cells
Species	Human
Predicted Molecular Mass	78.8 KDa
Formulation	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 6% Sucrose, 4% Mannitol, 0.05% Tween 80, pH8.0.
Shipping	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	

SDS-PAGE image



Background

Alternative Names	cGMP-Dependent Protein Kinase 1; cGK 1; cGK1; cGMP-Dependent Protein Kinase I; cGKI; PRKG1; PRKG1B; PRKGR1A; PRKGR1B
Background	cGMP-Dependent Protein Kinase 1 (PRKG1) belongs to the protein kinase superfamily and AGC Ser/Thr protein kinase family. PRKG1 contains one AGC-



kinase C-terminal domain, two cyclic nucleotide-binding domains, and one protein kinase domain. PRKG1 is mainly expressed in the lung and placenta. PRKG1 acts as a key mediator of the nitric oxide (NO)/cGMP signaling pathway. PRKG1 can phosphorylate many proteins that regulate platelet activation and adhesion, smooth muscle contraction, cardiac function, gene expression, feedback of the NO-signaling pathway, and other processes involved in several aspects of the CNS like axon guidance, hippocampal and cerebellar learning, circadian rhythm, and nociception.

Note

For Research Use Only, Not for Diagnostic Use.