


Summary

Name	Proteoglycan 3/PRG3
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Proteoglycan 3 is produced by our Mammalian expression system and the target gene encoding Leu18-Phe225 is expressed with a 6His tag at the C-terminus.
Accession #	AAI13412.1
Host	Human Cells
Species	Human
Predicted Molecular Mass	24.6 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human PRG3 (C-6His)
Catalog #: PHH1393



Background

Alternative Names

Proteoglycan 3; Eosinophil Major Basic Protein Homolog; Prepro-Major Basic Protein Homolog; Prepro-MBPH; PRG3; MBPH

Background

Proteoglycan 3, also known as Eosinophil major basic protein homolog, Prepro-major basic protein homolog, PRG3 and MBPH, contains one C-type lectin domain. Proteoglycans are a major component of the animal extracellular matrix. PRG3 localizes to the eosinophil secondary granule and is expressed in bone marrow, not detected in placenta. PRG3 has similar cytotoxic and cyto stimulatory activities to PRG2/MBP. In vitro, PRG3 can stimulate neutrophil superoxide production and IL8 release, histamine and leukotriene C4 release from basophils.

Note

For Research Use Only , Not for Diagnostic Use.