Catalog #: PHH1481



## **Summary**

Name	Semaphorin 3C/SEMA3C
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction Accession #	Recombinant Human Semaphorin 3C is produced by our Mammalian expression system and the target gene encoding Gly21-Ser738 is expressed with a 6His tag at the C-terminus. Q99985
Host	Human Cells
Species	Human
Predicted Molecular Mass	82.44 KDa
Formulation	Supplied as a 0.2 µm filtered solution of 20mM NaH2POsub//u003E4/sub//u003E, 150mM NaCl, 0.1M Arginine, 0.1M Glu, 10% Glycerol, 0.01% Tween20, 5% Trehalose, pH 7.4.
Shipping	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	-

## **SDS-PAGE** image

kDa	MK	R
120 90		-
60		
40	—	
30		
20	-	1000

## Background

Alternative Names	Semaphorin-3C; Semaphorin-E; Sema E; SEMA3C; SEMAE	
Background	Semaphorin 3C is one of six Class 3 secreted semaphorins, which are potent	



chemorepellents, that play a role in axon and/or vascular guidance during development can be up-regulated in the process of tumor progression. Semaphorin 3C consists of a 20 amino acids signal sequence, a N-terminal Sema domain, a cysteine knot, a furine-type cleavage site, an Ig-like domain, and a C-terminal basic domain. Semaphorin 3C expressed highly in the heart, colon, skeletal muscle, small intestine, testis, ovary and prostate, expressed lowly in the other organs, including the brain.

## **Note** For Research Use Only , Not for Diagnostic Use.