

Summary

Name Serpin B3/SCCA-1

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Serine Protease Inhibitor-clade B3 is produced by our

Mammalian expression system and the target gene encoding Met1-Pro390 is

expressed with a 6His tag at the C-terminus.

Accession # P29508

Host Human Cells

Species Human

Predicted Molecular Mass 45.96 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 0.02% Tween80,4%

Mannitol, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human Serpin B3 (C-6His) Enkilife Catalog #: PHH1505

Alternative Names Serpin B3; Protein T4-A; Squamous cell carcinoma antigen 1; SCCA-1;serine (or

cysteine) proteinase inhibitor, clade B (ovalbumin), member 3; serpin peptidase inhibitor, clade B (ovalbumin), member 3; Squamous cell carcinoma antigen 1; T4-

A;SCCA1

Background Serpin B3, also known as squamous cell carcinoma antigen-1 (SCCA-1), is a

member of the serpin superfamily of serine protease inhibitors. Serpin B3 belongs to the subgroup ovalbumin-related serpins which are involved in the regulation of apoptosis, inflammation, angiogenesis and embryogenesis. It may act as a papain-like cysteine protease inhibitor to modulate the host immune response against tumor cells. It also functions as an inhibitor of UV-induced apoptosis via

suppression of the activity of c-Jun NH(2)-terminal kinase (JNK1).

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838