Product Name: Recombinant Human TFF2 (C-6His)

Catalog #: PHH1618



Summary

TFF2/Trefoil factor 2 Name

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Human Trefoil Factor 2 is produced by our Mammalian

expression system and the target gene encoding Glu24-Tyr129 is expressed

with a 6His tag at the C-terminus.

Accession # Q03403

Host **Human Cells**

Species Human

Predicted Molecular Mass 13 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 Stability&Storage


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human TFF2 (C-6His)

Catalog #: PHH1618



Background

Alternative Names Trefoil Factor 2; Spasmolysin; Spasmolytic Polypeptide; SP; TFF2; SML1

Background Trefoil Factor 2 (TFF2) is a member of the trefoil family and contains two P-type

(trefoil) domains. Members of this family are characterized by having at least one copy of the trefoil motif, a 40-amino acid domain that contains three conserved disulfides. TFF2 is a secreted protein and specifically expressed in the stomach. TFF2 inhibits gastrointestinal motility and gastric acid secretion. TFF2 could function as a structural component of gastric mucus, possibly by stabilizing glycoproteins in the mucus gel through interactions with carbohydrate side chains.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838