
Catalog #: PHH1751



Summary

Name	TXNDC15/Thioredoxin domain-containing protein 15
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Thioredoxin Domain-Containing Protein 15 is produced by our Mammalian expression system and the target gene encoding Val33- Ser321 is expressed with a 6His tag at the C-terminus.
Accession #	Q96J42
Host	Human Cells
Species	Human
Predicted Molecular Mass	32.5 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human TXNDC15 (C-6His) Catalog #: PHH1751



Alternative NamesThioredoxin domain-containing protein 15;C5orf14;UNQ335/PRO534BackgroundThioredoxin domain-containing protein 15(TXNDC15) is a single-pass type I
membrane protein. Mature Human TXNDC15 consists of a 289 amino acid (aa)
extracellular region (ECD) with one thioredoxin domain, a 21 aa transmembrane
domain, and a 18 aa cytoplasmic region. It has 2 isoforms produced by alternative
splicing. Thioredoxins comprise a family of small proteins that, by catalyzing the
oxidation of disulfide bonds, participate in redox reactions throughout the cell.
Proteins that contain thioredoxins, but generally function as disulfide isomerases that
enzymatically rearrange disulfide bonds found in various proteins.

Note

For Research Use Only, Not for Diagnostic Use.