


Summary

Name	Uroplakin-2/UPK2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Uroplakin-2 is produced by our Mammalian expression system and the target gene encoding Asp26-Gly155 is expressed with a 6His tag at the C-terminus.
Accession #	O00526
Host	Human Cells
Species	Human
Predicted Molecular Mass	14.84 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human UPK2 (C-6His)
Catalog #: PHH1796



Alternative Names

Uroplakin-2; UP2; Uroplakin II; UPII; UPK2

Background

Uroplakin-2 is a single-pass type I membrane protein that belongs to the uroplakin-2 family. Uroplakin-2 is a component of the asymmetric unit membrane (AUM) and expressed in the ureter, a highly specialized biomembrane elaborated by terminally differentiated urothelial cells. Uroplakin-2 forms heterodimer with UPK1A that is necessary for exiting out of the endoplasmic reticulum (ER). Uroplakin-2 may play an important role in regulating the assembly of the AUM. AUM is believed to strengthen the urothelium by preventing cell rupture during bladder distention.

Note

For Research Use Only , Not for Diagnostic Use.