


Summary

Name	VSIG2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human V-Set and Immunoglobulin Domain-Containing Protein 2 is produced by our Mammalian expression system and the target gene encoding Val24-Ala243 is expressed with a 6His tag at the C-terminus.
Accession #	Q96IQ7
Host	Human Cells
Species	Human
Predicted Molecular Mass	24.2 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human VSIG2 (C-6His)
Catalog #: PHH1824



Alternative Names

V-Set and Immunoglobulin Domain-Containing Protein 2; Cortical Thymocyte-Like Protein; CT-Like Protein; VSIG2; CTH; CTXL

Background

V-Set and Immunoglobulin Domain-Containing Protein 2 (VSIG2) is presumably a 50-60 kDa single-pass type I transmembrane (glyco)protein which contains one Ig-like C2-type (immunoglobulin-like) domain and one Ig-like V-type (immunoglobulin-like) domain. VSIG2 is highly expressed in the stomach, colon, prostate, trachea and thyroid glands and weakly in bladder and lung. V-set domains are Ig-like domains resembling the antibody variable domain. V-set domains are found in diverse protein families, including immunoglobulin light and heavy chains, in several T-cell receptors such as CD2 (Cluster of Differentiation 2), CD4, CD80, and CD86, in myelin membrane adhesion molecules, in junction adhesion molecules (JAM), in tyrosine-protein kinase receptors, and in the programmed cell death protein 1 (PD1). It shows expression in stomach and prostate by Northern blot, and likely participates in cell adhesion. Human VSIG2 precursor is 327 amino acids in length.

Note

For Research Use Only , Not for Diagnostic Use.