


Summary

Name	MMP-8/Matrix metalloproteinase-8
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Neutrophil Collagenase is produced by our Mammalian expression system and the target gene encoding Phe21-Gly467 is expressed with a 10His tag at the C-terminus. The proenzyme needs to be activated by APMA for an activated form.
Accession #	P22894
Host	Human Cells
Species	Human
Predicted Molecular Mass	52.8 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human MMP-8 (C-10His)
Catalog #: PHH1905



Alternative Names

Neutrophil collagenase; Matrix metalloproteinase-8; MMP-8; PMNL collagenase; PMNL-CL; MMP8; CLG1

Background

Matrix metalloproteinases (MMPs) are a family of zinc and calcium dependent endopeptidases with the combined ability to degrade all the components of the extracellular matrix. MMP8(neutrophil collagenase) is expressed in neutrophils, where it is stored in specific granules. MMP8 release from the neutrophils is stimulated by various factors such as interleukins 1 and 8, TNF- α and GM-CSF. MMP8 is capable of cleaving types I, II and III triple-helical collagen, gelatin peptides, fibronectin, proteoglycans, aggrecan, serpins, β -casein and peptides such as angiotensin and substance P. In addition to its function in phagocytosis, MMP8 has a high capacity for infiltrating connective tissue, and is implicated in the breakdown of the extracellular matrix in diseases such as rheumatoid arthritis. MMP8 is heavily glycosylated.

Note

For Research Use Only , Not for Diagnostic Use.