


## Summary

<b>Name</b>	VEGFB/Vascular Endothelial Growth Factor B
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/ $\mu$ g as determined by LAL test.
<b>Construction</b>	Recombinant Human Vascular Endothelial Growth Factor B is produced by our Mammalian expression system and the target gene encoding Pro22-Ala207 is expressed with a human IgG1 Fc tag at the N-terminus.
<b>Accession #</b>	P49765
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	45.7 KDa
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution of 20 mM Glycine, 6% Sucrose, 5% Mannitol, 0.05% Tween80, pH3.0.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at $\leq -70^{\circ}\text{C}$ , stable for 6 months after receipt. Store at $\leq -70^{\circ}\text{C}$ , stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\mu$ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Human VEGFB (N-Fc)**  
**Catalog #: PHH1980**



---

**Alternative Names**

Vascular endothelial growth factor B;VEGF-B;VEGF-related factor;VRF

**Background**

VEGFB, as known as VRF, is a member of the VEGF family of growth factors that share structural and functional similarity. By alternative splicing, two isoforms of mature VEGF-B containing 167 or 186 amino acid (aa) residues exist. VEGF-B is expressed in most tissues, especially in heart, skeletal muscle and pancreas. The two VEGF-B isoforms have identical amino-terminal cysteine-knot VEGF homology domains but the carboxyl end of VEGF-B167 differs from that of VEGF-B186 by the presence of a highly basic cysteine-rich heparin binding domain. VEGF-B167 and a proteolytically processed form of VEGF-B186 also bind neuropilin-1, a type I transmembrane receptor for semaphorins/collapsins, ligands involved in neuron guidance.

**Note**

For Research Use Only , Not for Diagnostic Use.