Product Name: Recombinant Human PDYN (C-6His)

Catalog #: PHH1993



Summary

Name Proenkephalin-B/PDYN

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Proenkephalin-B is produced by our Mammalian

expression system and the target gene encoding Asp21-Ala254 is expressed

with a 6His tag at the C-terminus.

Accession # P01213

Host Human Cells

Species Human

Predicted Molecular Mass 27.3 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 4mM HCl.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human PDYN (C-6His) Catalog #: PHH1993

C EnkiLife

Alternative Names ADCA; PENKB; SCA23; PDYN; Dynorphin B; Big Dyn

Background Proenkephalin-B(PDYN), belongs to the opioid neuropeptide precursor family. The

N-terminal domain contains 6 conserved cysteines thought to be involved in disulfide bonding and/or processing. Leu-enkephalins, which is a type of Proenkephalin-B, compete with and mimic the effects of opiate drugs. They play a role in a number of physiologic functions, including pain perception and responses to stress. Dynorphin peptides differentially regulate the kappa opioid receptor. Dynorphin A has a typical opiod activity, it is 700 times more potent than Leu-

enkephalin.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838