
Product Name: Recombinant Human Cyclophilin B (C-6His) Catalog #: PHH1995



## Summary

Name	Cyclophilin B/PPIB/CYP-S1/CYPB/HEL-S-39/OI9/SCYLP
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Peptidyl-prolyl cis-trans Isomerase B is produced by our Mammalian expression system and the target gene encoding Asp34-Ala212 is expressed with a 6His tag at the C-terminus.
Accession #	P23284
Host	Human Cells
Species	Human
Predicted Molecular Mass	20.9 KDa
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 6% Sucrose, 4% Mannitol, 50mM NaCl, 0.05% Tween 80, pH6.0.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

# SDS-PAGE image



## Background



#### Alternative Names

CYP-S1; CYPB; HEL-S-39; OI9; SCYLP

Background

Cyclophilin B (SCYLP, CyPB and peptidyl-prolyl cis-trans isomerase B) is a 24 kDa glycoprotein member of the B subfamily of the cyclophilin-type PPlase family of molecules. It is both secreted and retained in the ER. When secreted, Cyclophilin B mediates chemotaxis and T cell adhesion to fibronectin. This is likely due to its prolyl cis/trans isomerase activity. Intracellularly, Cyclophilin B appears to serve as a molecular chaperone for molecules destined for secretion. It does so via stabilization, and facilitating the activity of additional chaperones.

#### Note

For Research Use Only, Not for Diagnostic Use.