Product Name: Recombinant Human CD316 (C-Fc)

Catalog #: PHH2192



Summary

Name CD316/IGSF8/Immunoglobulin Superfamily Member 8

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Immunoglobulin Superfamily Member 8 is produced by

our Mammalian expression system and the target gene encoding Arg28-

Thr579 is expressed with a human IgG1 Fc tag at the C-terminus.

Accession # Q969P0

Host Human Cells

Species Human

Predicted Molecular Mass 85.7 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human CD316 (C-Fc) Catalog #: PHH2192



Alternative Names Immunoglobulin Superfamily Member 8; IgSF8; CD81 Partner 3; Glu-Trp-Ile EWI

Motif-Containing Protein 2; EWI-2; Keratinocytes-Associated Transmembrane Protein 4; KCT-4; LIR-D1; Prostaglandin Regulatory-Like Protein; PGRL; CD316;

IGSF8; CD81P3; EWI2; KCT4

Background Immunoglobulin Superfamily Member 8 (IGSF8) is a single-pass membrane

protein. IGSF8 contains four Ig-like C2 type domains. The Ig-like C2-type domains 3 and 4 are required for interactions with CD81. IGSF8 may regulate proliferation and differentiation of keratinocytes. IGSF8 may participate in the regulation of neurite outgrowth and maintenance of the neural network in the adult brain. It also

may play a role on integrin-dependent morphology and motility functions.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838