


Summary

Name	Frizzled-8
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Frizzled-8 is produced by our Mammalian expression system and the target gene encoding Ala28-Pro172 is expressed with a 6His tag at the C-terminus.
Accession #	Q9H461
Host	Human Cells
Species	Human
Predicted Molecular Mass	17.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human Frizzled-8 (C-6His)
Catalog #: PHH2244



Alternative Names

frizzled 8; frizzled family receptor 8; Frizzled8; Frizzled-8; FZ-8; FZD8; hFZ8

Background

Frizzled-8 is one of at least ten seven-transmembrane (7TM) glycoproteins of the Frizzled family of Wnt receptors. Frizzled proteins are thought to be G-protein-coupled. Wnt engagement, with low density lipoprotein receptor-related proteins LRP-5 or LRP-6 acting as co-receptors, stabilizes beta-catenin and promotes gene transcription that is important in development and tissue maintenance. Component of the Wnt-Fzd-LRP5-LRP6 complex that triggers beta-catenin signaling through inducing aggregation of receptor-ligand complexes into ribosome-sized signalosomes. The beta-catenin canonical signaling pathway leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. These ligands bind the extracellular CRD of Frizzled-8, blocking Wnt binding. The recombinant Frizzled-8 CRD has also been used to block Wnt signaling and inhibit growth of teratocarcinomas.

Note

For Research Use Only , Not for Diagnostic Use.