
Product Name: Recombinant Human GIPR N-ECD (C-6His) Catalog #: PHH2469

# (C-6HIS) EnkiLife

## Summary

Name	GIPR
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Gastric inhibitory polypeptide receptor is produced by our Mammalian expression system and the target gene encoding Arg22-Gln138 is expressed with a 6His tag at C-terminus.
Accession #	P48546
Host	Human Cells
Species	Human
Predicted Molecular Mass	14.3 KDa
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### **SDS-PAGE** image



#### Background

Product Name: Recombinant Human GIPR N-ECD (C-6His) Catalog #: PHH2469

Alternative NamesGastric inhibitory polypeptide receptor; GIP-R; Glucose-dependent insulinotropic<br/>polypeptide receptor; GIPRBackgroundGIP receptor (GIPR) belongs to the G-protein coupled receptor family, activating<br/>adenylate cyclase and increasing levels of intracellular cyclic adenosine<br/>monophosphate (cAMP) in pancreatic b cells, thereby stimulating insulin section<br/>glucosedependently. New discoveries of GIP receptor (GIPR) biology in adipose<br/>tissue, as well as findings that co-agonists for the glucagon-like peptide-1 receptor<br/>(GLP-1R) and GIPR induce greater weight loss than that seen with GLP-1R agonists<br/>alone, has led to continued interest in manipulating GIPR activity for the treatment<br/>of obesity/type 2 diabetes mellitus (T2DM).

#### Note

For Research Use Only, Not for Diagnostic Use.