


## Summary

<b>Name</b>	Angiotensin-related protein 4/ANGPTL4
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Mouse Angiotensin Like Protein 4 is produced by our Mammalian expression system and the target gene encoding Lys167-Ser410 is expressed with a human IgG1 Fc tag at the C-terminus.
<b>Accession #</b>	Q9Z1P8
<b>Host</b>	Human Cells
<b>Species</b>	Mouse
<b>Predicted Molecular Mass</b>	54.6 KDa
<b>Formulation</b>	Supplied as a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	

## SDS-PAGE image



## Background

<b>Alternative Names</b>	Angiotensin-related protein 4;425O18-1;Angiotensin-like protein 4;Fasting-induced adipose factor;Hepatic fibrinogen/angiotensin-related protein;HFARP;Secreted protein Bk89;Angptl4;Farp; Fiaf; Ng27
--------------------------	--

**Product Name: Recombinant Mouse ANGPTL4 (C-Fc)**  
**Catalog #: PHM0064**



---

**Background**

Angiopoietin-related protein 4 (ANGPTL4) is a secreted protein and contains 1 fibrinogen C-terminal domain. The protein may act as a regulator of angiogenesis and modulate tumorigenesis. It inhibits proliferation, migration, and tubule formation of endothelial cells and reduces vascular leakage. ANGPTL4 may exert a protective function on endothelial cells through an endocrine action. It is directly involved in regulating glucose homeostasis, lipid metabolism, and insulin sensitivity (By similarity). In response to hypoxia, the unprocessed form of the protein accumulates in the subendothelial extracellular matrix (ECM). The matrix-associated and immobilized unprocessed form limits the formation of actin stress fibers and focal contacts in the adhering endothelial cells and inhibits their adhesion. It also decreases motility of endothelial cells and inhibits the sprouting and tube formation.

**Note**

For Research Use Only , Not for Diagnostic Use.