


Summary

Name	ANGPTL3/Angiopoietin-like Protein 3
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Angiopoietin-related Protein 3 is produced by our Mammalian expression system and the target gene encoding Ser17-Thr206 is expressed with a 6His tag at the C-terminus.
Accession #	Q9R182
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	22.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse ANGPTL3 (C-6His)
Catalog #: PHM0066



Alternative Names

Angiopoietin-related Protein 3; Angiopoietin-like protein 3; Angptl3

Background

Angiopoietin-like Protein 3 (ANGPTL3) is a secreted glycoprotein that is structurally related to the angiopoietins. Mature mouse ANGPTL3 contains an N-terminal coiled coil domain and a C-terminal fibrinogen-like domain. ANGPTL3 is expressed in the liver from early in development through adulthood. ANGPTL3 directly inhibits lipoprotein lipase (LPL) and endothelial lipase (EL), enzymes responsible for hydrolyzing circulating triglycerides and HDL phospholipids. This activity requires a putative heparin-binding motif which is N-terminal to the coiled coil domain. Proteolytic removal of the fibrinogen-like domain from the N-terminal fragment serves to activate ANGPTL3 and increase its ability to inhibit LPL in vitro and function in vivo. ANGPTL3 promotes an increase in circulating triglyceride levels without altering VLDL or HDL secretion or uptake. ANGPTL3 expression in vivo is up-regulated by LXR agonists and down-regulated by insulin, leptin, and agonists of TR β or PPAR β . ANGPTL3, secreted by fetal liver cells, also promotes the expansion of hematopoietic stem cells.

Note

For Research Use Only , Not for Diagnostic Use.