


Summary

Name	COL18A1/Collagen alpha-1(XVIII) Chain/Endostatin
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/ μ g as determined by LAL test.
Construction	Recombinant Mouse Collagen Alpha-1(XVIII) Chain is produced by our Mammalian expression system and the target gene encoding His1591-Lys1774 is expressed with a 6His tag at the C-terminus.
Accession #	P39061
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	21.2 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse Endostatin (C-6His)
Catalog #: PHM0432

Alternative Names	antiangiogenic agent;COL18A1;collagen alpha-1(XVIII)chain; collagen;type XVIII;Endostatin
Background	Endostatin, an endogenous non-glycosylated inhibitor of endothelial cell proliferation and angiogenesis. It is produced and/or trimmed by metalloproteinases such as MMP-2 and MMP-9, and cathepsins S, B and L. The N-terminal ~27 aa of Endostatin appear to contain the majority of its activity. This region contains zinc binding sites that are thought to be critical for its anti-endothelial and anti-tumor effects, as well as multiple cleavage sites that, when used, can modify its activity. Mouse Endostatin shares 96% aa sequence identity with rat and 85-87% with human, bovine and equine Endostatin. It is predominantly expressed in liver, kidney, lung, skeletal muscle and testis. Endostatin inhibits endothelial cell growth by inducing cell cycle arrest in G1 phase and initiating apoptosis. It is also thought to down-regulate angiogenesis by blocking VEGF-induced endothelial cell migration. Endostatin may also be involved with down-regulation of angiogenesis after establishment of placental circulation in the pregnant uterus.

Note

For Research Use Only , Not for Diagnostic Use.