


## Summary

<b>Name</b>	CX3CL1/Fractalkine
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Mouse C-X3-C motif Chemokine 1 is produced by our Mammalian expression system and the target gene encoding Gln25-Arg337 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	O35188
<b>Host</b>	Human Cells
<b>Species</b>	Mouse
<b>Predicted Molecular Mass</b>	34.3 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Mouse CX3CL1 (C-6His)**  
**Catalog #: PHM0462**



---

**Alternative Names**

Fractalkine; C-X3-C motif chemokine 1; CX3C membrane-anchored chemokine; Neurotactin; Small-inducible cytokine D1; Cx3c; Fkn; Scyd1; CXC3; CXC3C; ABCD-3; SCYD1; C3Xkine; NTN; NTT

**Background**

Fractalkine(CX3CL1) is a single-pass type I membrane protein and belongs to the intercrine delta family. It consists of an extracellular NH<sub>2</sub>-terminal domain, a mucin-like stalk, a transmembrane  $\alpha$  helix, and a short cytoplasmic tail. CX3CL1 exists in two forms: as a membrane-anchored or as a shed 80-95K glycoprotein. Soluble CX3CL1 is generated by limited proteolysis on the cell surface, and a disintegrin and metalloproteinase 10 (ADAM10) and ADAM17/tumor necrosis factor- $\alpha$ -converting enzyme (ADAM17/TACE) participate in this shedding. It has been suggested that ADAM10 acts in the constitutive shedding, and ADAM17 acts in response to cell activation. The protein may play a role in regulating leukocyte adhesion and migration processes at the endothelium.

**Note**

For Research Use Only , Not for Diagnostic Use.