


Summary

Name	CXCL1/C-X-C motif chemokine 1/KC/GRO alpha/CINC1/MGSA-alpha
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse C-X-C Motif Chemokine 1 is produced by our Mammalian expression system and the target gene encoding Arg20-Lys96 is expressed with a 6His tag at the C-terminus.
Accession #	P12850
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	9.4 kDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Mouse CXCL1 (C-6His)
Catalog #: PHM0463



Background

Alternative Names

Growth-regulated alpha protein;C-X-C motif chemokine 1;Platelet-derived growth factor-inducible protein KC;Secretory protein N51;Cxcl1;Gro; Gro1; Mgsa; Scyb1

Background

Growth-regulated alpha protein (CXCL1,KC), is a member of the alpha chemokine subfamily, was initially identified as an immediate early gene induced in mouse fibroblasts by platelet-derived growth factor. The N-terminal processed form KC(5-72) of the protein is produced by proteolytic cleavage after secretion from bone marrow stromal cells, and shows a highly enhanced hematopoietic activity. Mouse KC shows approximately 63% identity to that of mouse MIP-2. KC is also approximately 60% identical to the human GROs. It has been suggested that mouse KC and MIP-2 are the orthologs of the human GROs and rat CINC3s. Cxcl1 has chemotactic activity for neutrophils, and contributes to neutrophil activation during inflammation. Hematopoietic chemokine, in vitro, suppresses hematopoietic progenitor cell proliferation.

Note

For Research Use Only , Not for Diagnostic Use.