


## Summary

Name	Frizzled-1/Fz-1/Fzd1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Mouse Frizzled (Drosophila) Homolog 1 is produced by our Mammalian expression system and the target gene encoding Val69-His248 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	070421
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	46.8 KDa
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## **SDS-PAGE** image



## Background



Alternative Names	Frizzled-1; Fz-1; mFz1; Fzd1; Frizzled homolog 1
Background	Frizzled homolog 1 (FZD1), also known as Frizzled-1, a member of the G-protein- coupled receptor superfamily, mediates Wnt signaling by serving as a co-receptor with LRP-5 for Wnt ligands. The FZD1 protein contains a signal peptide, a cysteine- rich domain in the N-terminal extracellular region, 7 transmembrane domains, and a C-terminal PDZ domain-binding motif.

## Note

For Research Use Only, Not for Diagnostic Use.