


Summary

Name	IL-21R/Interleukin-21 Receptor/IL21R
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Interleukin-21 Receptor is produced by our Mammalian expression system and the target gene encoding Cys20-Pro236 is expressed with a 6His tag at the C-terminus.
Accession #	Q9JHX3
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	25.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse IL-21R (C-6His)
Catalog #: PHM0995



Alternative Names

Interleukin-21 receptor; IL-21 receptor; IL-21R; IL21r; Nilr

Background

Interleukin-21 receptor (IL-21R) is a type I transmembrane glycoprotein within the class I cytokine receptor family, type 4 subfamily. IL-21R is expressed mainly on B cells, NK cells, and activated T cells, but is also found on dendritic cells, alternatively activated macrophages, intestinal lamina propria fibroblasts and epithelial cells, and keratinocytes. Both IL-21 and IL-4 are necessary for efficient B cell IgG1 production and normal germinal center architecture. B cell IL-21 R engagement induces Blimp-1 (which mediates plasma cell differentiation), and is important for memory responses. IL-21R engagement on mouse NK cells enhances their cytotoxic activity and IFN- γ production. IL-21R engagement on CD8+ T cells aids control of viral infection and tumor growth; IL-21R is also necessary for sufficient numbers of regulatory T cells to combat chronic inflammation. IL-21R expression is often up-regulated in allergic skin inflammation, systemic lupus erythematosus and diffuse large B cell lymphoma (DLBCL).

Note

For Research Use Only , Not for Diagnostic Use.