# Product Name: Recombinant Mouse MARCO (N-8His) Catalog #: PHM1134



### **Summary**

Name MARCO

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Mouse Macrophage Receptor MARCO is produced by our

Mammalian expression system and the target gene encoding Gln70-Ser518 is

expressed with a 8His tag at the N-terminus.

Accession # Q60754

**Host** Human Cells

**Species** Mouse

Predicted Molecular Mass 46.2 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### **SDS-PAGE** image



## **Background**

# Product Name: Recombinant Mouse MARCO (N-8His) Catalog #: PHM1134



**Alternative Names** 

Macrophage receptor MARCO; Marco; Macrophage receptor with collagenous

structure

**Background** 

MARCO (macrophage receptor with collagenous structure) is an 80 kDa type II transmembrane glycoprotein that belongs to the class A scavenger receptor family. Mouse MARCO consists of a 48 amino acid (aa) cytoplasmic domain, a 21 aa transmembrane segment, and a 449 aa extracellular domain (ECD) that includes a stalk region, a collagenlike region, and one SRCR domain. MARCO is constitutively expressed on the surface of splenic and lymph node macrophages. MARCO binds LPS, lipoteichoic acid, and other determinants on Gram positive and Gram negative bacteria. It also binds modified LDL, CpG oligonucleotides, UGRP1, silica, and TiO2.

#### Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838