


Summary

| | |
|---------------------------------|--|
| Name | SLAMF4/Natural killer cell receptor 2B4/CD244 |
| Purity | Greater than 95% as determined by reducing SDS-PAGE |
| Endotoxin level | <1 EU/μg as determined by LAL test. |
| Construction | Recombinant Mouse Natural Killer Cell Receptor 2B4 is produced by our Mammalian expression system and the target gene encoding Gln20-Asn221 is expressed with a 6His tag at the C-terminus. |
| Accession # | Q07763 |
| Host | Human Cells |
| Species | Mouse |
| Predicted Molecular Mass | 23.5 KDa |
| Formulation | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. |
| Stability&Storage | Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles. |
| Reconstitution | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

SDS-PAGE image



Background

Product Name: Recombinant Mouse CD244 (C-6His)
Catalog #: PHM1194



Alternative Names

Natural killer cell receptor 2B4; NK cell type I receptor protein 2B4; NKR2B4; SLAM family member 4; SLAMF4; Signaling lymphocytic activation molecule 4; CD244

Background

Natural killer cell receptor 2B4 (2B4/CD244) is a 66 kDa type I transmembrane glycoprotein in the SLAM subgroup of the CD2 protein family. SLAM family proteins have an extracellular domain (ECD) with two or four Ig-like domains and at least two cytoplasmic immunoreceptor tyrosine-based switch motifs (ITSMs). 2B4 interacts with CD48, while other SLAM family proteins interact in a homophilic manner. The mouse 2B4 cDNA encodes a 397 amino acid (aa) precursor that includes a 19 aa signal sequence, a 207 aa ECD with one Ig-like V-type and one C2-type Ig-like domain, a 21 aa transmembrane segment, and a 150 aa cytoplasmic domain with four ITSMs. Within the ECD, mouse 2B4 shares 46% and 68% aa sequence identity with human and rat 2B4, respectively. 2B4/CD48 signaling cooperates with other receptor systems to either promote or inhibit NK and CD8+ T cell activation. The inhibitory activities are distinct from those of MHC I restricted inhibitory NK cell receptors. Ligation of 2B4 with antibodies or CD48 constructs can directly trigger inhibitory signaling or disrupt an inhibitory interaction, leading to cellular activation. 2B4 can also induce signaling through CD48.

Note

For Research Use Only , Not for Diagnostic Use.