


Product Name: Recombinant Mouse SEMA4A (C-6His)
Catalog #: PHM1482



Summary

Name	Semaphorin 4A/SEMA4A
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/ μ g as determined by LAL test.
Construction	Recombinant Mouse Semaphorin 4A is produced by our Mammalian expression system and the target gene encoding Thr33-His682 is expressed with a 6His tag at the C-terminus.
Accession #	Q62178
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	72.7 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of 20mM PB, 150mM NaCl, 1mM EDTA, 5% Trehalose, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse SEMA4A (C-6His)
Catalog #: PHM1482



Alternative Names

Semaphorin-4A;Semaphorin-B;Sema B;Sema4a;Semab; SemB

Background

Semaphorin-4A (SEMA4A) belongs to the semaphorin family which contains a Ig-like C2-type domain, a PSI domain and a Sema domain. SEMA4A is expressed from day 10 in the embryo, and low levels are found between days 10-12. SEMA4A is a cell surface receptor for PLXNB1, PLXNB2, PLXNB3 and PLXND1 that plays an important role in cell-cell signaling. It plays a role in priming antigen-specific T-cells, promotes differentiation of Th1 T-helper cells, and thereby contributes to adaptive immunity. SEMA4A promotes phosphorylation of TIMD2, inhibits angiogenesis, and promotes axon growth cone collapse, Inhibits axonal extension by providing local signals to specify territories inaccessible for growing axons.

Note

For Research Use Only , Not for Diagnostic Use.