


Summary

Name	TSLP R/Thymic Stromal Lymphopoietin Receptor/CRLF2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Thymic Stromal Lymphopoietin Protein Receptor is produced by our Mammalian expression system and the target gene encoding Ala20-Leu233 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	AAH23788.1
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	49.8 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Mouse TSLP R (C-Fc)
Catalog #: PHM1740



Background

Alternative Names

CRL2; CRLF2; CRL2 cytokine receptor; cytokine receptor-like factor 2; ILXR; IL-XR; P2RY8/CRLF2 fusion; Thymic stromal lymphopoietin protein receptor; Thymic stromal-derived lymphopoietin receptor; TSLP receptor; TSLPR

Background

The cytokine thymic stromal lymphopoietin receptor (TSLPR) is consisting of a common γ receptor-like chain (TSLPR- γ) and a common interleukin 7 (IL-7) α chain that belongs to the type 1 cytokine receptor family. Transfection of TSLPR cDNA result in only low affinity binding, while cotransfection of the IL-7 α chain cDNA shows high affinity binding. TSLP and TSLPR play a critical role in the initiation of allergic diseases in mice. The TSLP R cDNA encodes a transmembrane receptor containing 370 amino acids (aa) with two potential N-linked glycosylation sites and a cytoplasmic domain of 104 aa including a single tyrosine residue. TSLPR can mediate signaling of the signal transducer and activator of transcription 5 (Stat5) by TSLP. TSLP R is broadly expressed in the immune and hematopoietic cells, particularly in hematopoietic progenitors and myeloid cells.

Note

For Research Use Only , Not for Diagnostic Use.