
Catalog #: PHM2179



## **Summary**

Name	SECTM1A
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Mouse Secreted And Transmembrane Protein 1A is produced by our Mammalian expression system and the target gene encoding Gln28/xadThr165 is expressed with a 6His tag at the C-terminus.
Accession #	A2ABP9
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	16.2 KDa
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

# **SDS-PAGE** image



# Background

Product Name: Recombinant Mouse SECTM1A (C-6His) Catalog #: PHM2179



#### **Alternative Names**

SECTM1A; Sectm1a

**Background** SECTM1A (secreted and transmembrane 1A), is 192 amino acid (aa) protein, appears to share structural and functional characteristics with other SECTM1 proteins. Human SECTM1 can be found either found as an approximately 27 kDa intracellular type I transmembrane protein that shows a perinuclear, Golgi like staining pattern, or as a 20 kDa soluble, secreted form, and is produced by some myeloid cells and by thymic epithelia and fibroblasts. Stimulation with IFN gamma is often necessary to detect human SECTM1 expression, and it is thought to be an interferon early response gene. Mouse SECTM1A cDNA encodes a signal sequence, an extracellular domain with four potential N linked glycosylation sites, a transmembrane sequence, and a very short (approximately 6 aa) cytoplasmic sequence. SECTM1 proteins from human and mouse show species specific binding of CD7 and co stimulation of T cells, including enhancement of CD3 induced proliferation.

### Note

For Research Use Only, Not for Diagnostic Use.