## **Product Name: Recombinant Mouse CD7 (C-6His)**

Catalog #: PHM2224



#### **Summary**

Name CD7/Leu-9/T-Cell Antigen CD7/GP40/T-Cell Surface Antigen Leu-9/TP41

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <0.01 EU/μg as determined by LAL test.

Construction Recombinant Mouse T-Cell Antigen CD7 is produced by our Mammalian

expression system and the target gene encoding Gln24/xadPro150 is

expressed with a 6His tag at the C-terminus.

Accession # P50283

**Host** Human Cells

**Species** Mouse

Predicted Molecular Mass 15.1 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### SDS-PAGE image



### **Background**

# **Product Name: Recombinant Mouse CD7 (C-6His)**

**C** EnkiLife

Catalog #: PHM2224

Alternative Names T-Cell Antigen CD7; GP40; T-Cell Leukemia Antigen; T-Cell Surface Antigen Leu-9;

TP41; CD7

Background T-Cell Antigen CD7 is a single-pass type I membrane protein that that belongs to

the the immunoglobulin superfamily. Human CD7 is synthesized as a 240 amino acid precursor that contains a 25 amino acid signal sequence and a 215 amino acid mature chain with a Ig-like (immunoglobulin-like) domain. CD7 is normally expressed on all T-lymphocytes, NK-cells, pre-B lymphocytes and pleuripotent hematopoietic stem cells. CD7 plays an essential role in T-cell interactions, T-cell/B-cell interaction during early lymphoid development, T- and NK-cell activation and cytokine production. CD7 has been shown to interact with PIK3R1and SECTM1. However, the function of the CD7 protein in the immune system is still largely

unknown.

#### Note

For Research Use Only , Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838