


**Product Name: Recombinant Mouse GDF15 (N-8His-Flag)**  
**Catalog #: PHM2405**



## Summary

<b>Name</b>	GDF15/GDF-15
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Mouse Growth Differentiation Factor/xa015 is produced by our Mammalian expression system and the target gene encoding Ser189-Ala303 is expressed with a 8His, Flag tag at the N-terminus.
<b>Accession #</b>	Q9Z0J7
<b>Host</b>	Human cells
<b>Species</b>	Mouse
<b>Predicted Molecular Mass</b>	16.9 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 4mM HCl.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in 4mM HCl. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in 4mM HCl. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Mouse GDF15 (N-8His-Flag)**  
**Catalog #: PHM2405**



---

**Alternative Names**

Growth Differentiation Factor/xa015, Macrophage inhibitory cytokine 1, GDF-15, MIC-1, NAG-1, PLAB, PTGFB

**Background**

Growth Differentiation Factor 15 (GDF-15), also called Macrophage Inhibitory Cytokine 1 (MIC-1), is a divergent member of the TGF-beta superfamily. GDF15 can be secreted by a wide variety of cell types in response to a broad range of stressors. GDF-15 expression is dramatically upregulated during acute brain injury, cancer, cardiovascular disease, and inflammation, suggesting its potential value as a disease biomarker. GDF15 was shown to inhibit proliferation of primitive hematopoietic progenitors and introduced as a putative placental mediator of embryonic development. GDF15 has recently gained scientific and translational prominence with the discovery that its receptor is a GFRAL-RET heterodimer of which GFRAL is expressed solely in the hindbrain.

**Note**

For Research Use Only , Not for Diagnostic Use.