## **Product Name: Recombinant Rat CD38 (C-6His)**

Catalog #: PHR1883



#### **Summary**

Name CD38/ADP-ribosyl Cyclase 1/cyclic ADP-ribose Hydrolase 1

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Rat ADP-ribosyl Cyclase/cyclic ADP-ribose Hydrolase 1 is

produced by our Mammalian expression system and the target gene

encoding Trp45-Val303 is expressed with a 6His tag at the C-terminus.

Accession # Q64244

**Host** Human Cells

**Species** Rat

Predicted Molecular Mass 30.8 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### **SDS-PAGE** image



## **Background**

# **Product Name: Recombinant Rat CD38 (C-6His)**

Catalog #: PHR1883



Alternative Names ADP-ribosyl cyclase 1; cyclic ADP-ribose hydrolase;CD38;T10

Background CD38, also called ADP-ribosyl cyclase, is a Type II integral membrane protein with

301 amino acids in length that belongs to the ADP-ribosyl cyclase family. It synthesizes the second messagers cyclic ADP-ribose and nicotinate-adenine dinucleotide phosphate, the former a second messenger for glucose-induced insulin secretion. And also moonlights as a receptor in cells of the immune system. CD38 is expressed in B and T lymphocytes, osteoclasts, and in cardiac, pancreatic, liver and kidney cells. Through its production of cyclic ADP-ribose, CD38 modulates calcium-mediated signal transduction in many types of cells, including

neutrophils and pancreatic beta cells.

#### Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838