Summary

Name CD38/ADP-ribosyl Cyclase 1/cyclic ADP-ribose Hydrolase 1

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Cynomolgus ADP-ribosyl Cyclase/cyclic ADP-ribose Hydrolase 1

is produced by our Mammalian expression system and the target gene

encoding Leu44-Ile301 is expressed with a 6His tag at the C-terminus.

Accession # Q5VAN0

Host Human Cells
Species Cynomolgus

Predicted Molecular Mass 30.9 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH

8.0.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Alternative Names ADP-ribosyl cyclase 1; cyclic ADP-ribose hydrolase; CD38; T10

Background CD38, also called ADP-ribosyl cyclase, is a Type II integral membrane protein with

301 amino acids in length that belongs to the ADP-ribosyl cyclase family. It synthesizes the second messagers cyclic ADP-ribose and nicotinate-adenine dinucleotide phosphate, the former a second messenger for glucose-induced insulin secretion. And also moonlights as a receptor in cells of the immune system. CD38 is expressed in B and T lymphocytes, osteoclasts, and in cardiac, pancreatic, liver and kidney cells. Through its production of cyclic ADP-ribose, CD38 modulates calcium-mediated signal transduction in many types of cells, including

neutrophils and pancreatic beta cells.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838