Summary

Name CD73/5'-Nucleotidase/NT5E

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Cynomolgus 5-nucleotidase/CD73 is produced by our

Mammalian expression system and the target gene encoding Trp27-Lys547 is

expressed with a 6His tag at the C-terminus.

Accession # G7P3J1

Host Human Cells
Species Cynomolgus

Predicted Molecular Mass 58.8 KDa

Formulation Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 120mM NaCl, 4mM

CaCl2, 20% Glycerol, pH 7.5.

Shipping The product is shipped on dry ice/polar packs. Upon receipt, store it immediately

at the temperature listed below.


Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution

SDS-PAGE image



Background

Alternative Names 5-Nucleotidase; 5-NT; Ecto-5-Nucleotidase; CD73; NT5E; NT5; NTE

Background CD73, also known as Ecto-5-nucleotidase, is an ectoenzyme that is attached to the

cell membrane by a glycosyl phophatidylinositol anchor. The enzyme is expressed

by most cell types. The 5′-Nucleotidase activity of CD73 converts extracellular nucleoside-5′-monophosphates to nucleosides. CD73 is one of several enzymes responsible for the production of extracellular adenosine, a signaling molecule that is involved in responses to inflammation and tissue injury. CD73 hydrolyzes extracellular nucleotides into membrane permeable nucleosides.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838