


## Summary

<b>Name</b>	B7-1/CD80/T-lymphocyte Activation Antigen CD80
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Cynomolgus T-lymphocyte Activation Antigen CD80 is produced by our Mammalian expression system and the target gene encoding Val35-Asn242 is expressed with a human IgG1 Fc tag at the C-terminus.
<b>Accession #</b>	G7NXN7
<b>Host</b>	Human Cells
<b>Species</b>	Cynomolgus
<b>Predicted Molecular Mass</b>	51 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 50 mM Tris-HCl, 100 mM Glycine, pH 7.5.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



**Product Name: Recombinant Cynomolgus B7-1 (C-Fc)**  
**Catalog #: PHV1941**



---

## Background

**Alternative Names** T-lymphocyte activation antigen CD80; Activation B7-1 antigen; B7; CD80

**Background** Cynomolgous Cluster of Differentiation 80, also called B7-1, is a member of cell surface immunoglobulin superfamily. It is expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells. CD80 plays key, yet distinct roles in the activation of T cells. B7-1/CD80 and B7-2/CD86, together with their receptors CD28 and CTLA4, constitute one of the dominant co-stimulatory pathways that regulate T- and B- cell responses. CD80 is mostly expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells. Although both CTLA-4 and CD28 can bind to the same ligands, CTLA-4 binds to B7-1 and B7-2 with a 20-100 fold higher affinity than CD28 and is involved in the down-regulation of the immune response. CD80 is thus regarded as promising therapeutic targets for autoimmune diseases and various carcinomas.

## Note

For Research Use Only , Not for Diagnostic Use.