


Product Name: Recombinant Cynomolgus FcRn Heterodimer (C-Flag-6His)
Catalog #: PHV1967

Summary

Name	FcRn/FCGRT & B2M Heterodimer
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Cynomolgus Beta-2-microglobulin&IgG Receptor FcRn Large Subunit P51 is produced by our Mammalian expression system and the target gene encoding Ile21-Met119&Ala24-Ser297 is expressed with a Flag tag at the C-terminus, 6His tag at the C-terminus.
Accession #	Q8SPW0-1&Q8SPV9-1
Host	Human Cells
Species	Cynomolgus
Predicted Molecular Mass	12.6&31.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Cynomolgus FcRn Heterodimer (C-Flag-5His)
Catalog #: PHV1967

Alternative Names IgG receptor FcRn; Neonatal Fc receptor; FCRN

Background β -2-Microglobulin (B2M) is a secreted protein with 1 Ig-like C1-type (immunoglobulin-like) domain which belongs to the beta-2-microglobulin family. B2M component of major histocompatibility complex (MHC) class I molecules, involved in the presentation of peptide antigens to the immune system. Polymers of beta 2-microglobulin can be found in tissues from patients on long-term hemodialysis. B2M is a protein found on the surface of many cells and plentiful on the surface of white blood cells. Serum B2M concentration is increased in renal diseases, various malignant diseases and some inflammatory and autoimmune disorders. B2M may adopt the fibrillar configuration of amyloid in certain pathologic states. The capacity to assemble into amyloid fibrils is concentration dependent. B2M has been shown as a marker for monitoring inflammatory disease activity and it appears likely to have a destructive role in amyloidosis-related arthritis. B2M might be involved in the OA (osteoarthritis) pathogenesis. Defects in B2M are the cause of hypercatabolic hypoproteinemia. Affected individuals show marked reduction in serum concentrations of immunoglobulin and albumin, probably due to rapid degradation. B2M could be a potential therapeutic target in ovarian cancer.

Note

For Research Use Only , Not for Diagnostic Use.