
Catalog #: PHV1986



## Summary

Name	PD-L2/B7-DC/CD273/Programmed cell death ligand 2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Cynomolgus Monkey Programmed Cell Death Ligand 2 is produced by our Mammalian expression system and the target gene encoding Leu20-Pro219 is expressed with a human IgG1 Fc tag at the C- terminus.
Accession #	A4GW30
Host	Human Cells
Species	Cynomolgus
Predicted Molecular Mass	49.5 KDa
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than $100\mu g/ml$ . Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than $100\mu g/ml$ . Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## **SDS-PAGE** image



## Background

Product Name: Recombinant Cynomolgus PD-L2 (C-Fc) Catalog #: PHV1986

Alternative NamesProgrammed Cell Death 1 Ligand 2; PD-1 Ligand 2; PD-L2; PDCD1 Ligand 2;<br/>Programmed Death Ligand 2; Butyrophilin B7-DC; B7-DC; CD273; PDCD1LG2;<br/>B7DC; CD273; PDCD1L2; PDL2BackgroundProgrammed Cell Death 1 Ligand 2 (PDCD1LG2) is a member of the BTN/MOG<br/>family. PDCD1LG2 contains one Ig-like C2-type domain and one Ig-like V-type<br/>domain. PDCD1LG2 is highly expressed in the heart, placenta, pancreas, lung and<br/>liver; it is weakly expressed in the spleen, lymph nodes, and thymus. PDCD1LG2 is<br/>involved in the costimulatory signal, essential for T-cell proliferation and IFNG<br/>production in a PDCD1-independent manner. PDCD1LG2 interaction with PDCD1<br/>inhibits T-cell proliferation by blocking cell cycle progression and cytokine<br/>production.

EnkiLife

## Note

For Research Use Only, Not for Diagnostic Use.