


Summary

Name	CD79B/B29
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Rhesus Macaque CD79B is produced by our Mammalian expression system and the target gene encoding Ala30-Asp161 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	H9ZFT8
Host	Human Cells
Species	Rhesus Macaque
Predicted Molecular Mass	42.2 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Rhesus Macaque CD79B (C-Fc)
Catalog #: PHV2184



Alternative Names

B-Cell Antigen Receptor Complex-Associated Protein Beta Chain; B-Cell-Specific Glycoprotein B29; Ig-Beta; Immunoglobulin-Associated B29 Protein; CD79b; CD79B; B29; IGB

Background

CD79B is a single-pass type I membrane protein. CD79B contains one Ig-like V-type domain and one ITAM domain. CD79B is required in cooperation with CD79A for initiation of the signal transduction cascade activated by the B-cell antigen receptor complex (BCR), which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79B enhances phosphorylation of CD79A, possibly by recruiting kinases that phosphorylate CD79A or by recruiting proteins that bind to CD79A and protect it from dephosphorylation.

Note

For Research Use Only , Not for Diagnostic Use.