## **Summary**

Name TREM-2/Triggering Receptor Expressed On Myeloid 2

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Cynomolgus Triggering receptor expressed on myeloid cells 2 is

produced by our Mammalian expression system and the target gene

encoding His19-Ser174 is expressed with a 6His tag at the C-terminus.

Accession # XP 005553122.1

Host Human Cells
Species Cynomolgus

Predicted Molecular Mass 18.3 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### **SDS-PAGE** image



# **Background**

# EnkiLife Product Name: Recombinant Cynomolgus TREM-2 (C-6His) Catalog #: PHV2392

**Alternative Names** 

Ig-like domain-containing protein; TREM2; Triggering receptor expressed on

myeloid cells 2

**Background** 

TREM2 is a cell surface receptor of the immunoglobulin superfamily. TREM2 is a type-1 transmembrane protein that shuttles to the plasma membrane where it exerts its cell autonomous biological functions. TREM2 undergoes regulated intramembrane proteolysis (RIP). TREM2 is preferentially expressed in microglia and is functionally required for migration, cytokine release, phagocytosis, lipid sensing, ApoE binding, shielding of amyloid plaques, and microglia proliferation in the brian. Most of the functionally investigated mutations are located within the

Ig-like domain of TREM2.

#### Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838