


Product Name: Recombinant *Cavia porcellus* CTLA-4 (C-6His)
Catalog #: PPV0458



Summary

Name	CTLA-4/CD152/Cytotoxic T-lymphocyte Protein 4
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant <i>Cavia Porcellus</i> Cytotoxic T-lymphocyte Protein 4 is produced by our Yeast expression system and the target gene encoding Ala37-Asp161 is expressed with a 6His tag at the C-terminus.
Accession #	H0VUB1
Host	<i>P.Pichia</i>
Species	<i>Cavia porcellus</i>
Predicted Molecular Mass	16-35 KDa
Formulation	Supplied as a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	

SDS-PAGE image



Background

Alternative Names Cytotoxic T-lymphocyte protein 4; Cytotoxic T-lymphocyte-associated antigen 4; CTLA-4; CD152; CTLA4

Background Cytotoxic Tlymphocyte 4(CTLA-4,CD152), is a type I transmembrane T cell inhibitory molecule that is a member of the Ig superfamily.CD28 and CTLA-4,

Product Name: Recombinant Cavia porcellus CTLA-4 (C-6His)
Catalog #: PPV0458



together with their ligands, B7-1 and B7-2, constitute one of the dominant costimulatory pathways that regulate T and B cell responses. CD28 and CTLA-4 are structurally homologous molecules that are members of the immunoglobulin (Ig) gene superfamily. CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Intracellular CTLA4 is also found in regulatory T Cells and may play an important role in their functions. Tcell activation through the Tcell receptor and CD28 leads to increased expression of CTLA4.

Note

For Research Use Only , Not for Diagnostic Use.