

Catalog #: AMM04164



Summary

Acetyl Lysine(10B10)Mouse Monoclonal Antibody **Production Name**

Description Mouse Monoclonal Antibody

Host Mouse **Application** WB,IHC

Reactivity Species independent

Performance

Conjugation Unconjugated Modification Acetyl Antibody

Isotype IgG

Clonality Monoclonal

Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

PBS, pH 7.4, containing 0.5%BSA, 0.02% New type preservative N as Preservative and Buffer

50% Glycerol.

Purification Affinity purification

Immunogen

Gene Name

Alternative Names

Gene ID

SwissProt ID .Purified Protein

Application

Dilution Ratio WB 1:1000-2000 IHC 1:200-500

Molecular Weight

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Antibody

Catalog #: AMM04164




Background


Acetylation of lysine, like phosphorylation of serine, threonine or tyrosine, is an important reversible modification controlling protein activity. The conserved amino-terminal domains of the four core histones (H2A, H2B, H3, and H4) contain lysines that are acetylated by histone acetyltransferases (HATs) and deacetylated by histone deacetylases (HDACs). Signaling resulting in acetylation/deacetylation of histones, transcription factors, and other proteins affects a diverse array of cellular processes including chromatin structure and gene activity, cell growth, differentiation, and apoptosis. Recent proteomic surveys suggest that acetylation of lysine residues may be a widespread and important form of posttranslational protein modification that affects thousands of proteins involved in control of cell cycle and metabolism, longevity, actin polymerization, and nuclear transport. The regulation of protein acetylation status is impaired in cancer and polyglutamine diseases, and HDACs have become promising targets for anti-cancer drugs currently in development.

Research Area

Image Data



Western blot analysis of 1) Mouse Brain Tissue, 2) Hela, 3) Hela+TSA Treated using Acetyl Lysine Monoclonal Antibody.



Immunohistochemical analysis of paraffin-embedded Human Breast Carcinoma using Acetyl Lysine Monoclonal Antibody.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838



Catalog #: AMM04164



Note

For research use only.

Tel: 0086-27-87002838