

Summary

| Production Name | p38 Rabbit Monoclonal Antibody | |
|-----------------|--|--|
| Description | Recombinant Rabbit Monoclonal antibody | |
| Host | Rabbit | |
| Application | WB,IP | |
| Reactivity | Human | |

Performance

| Conjugation | Unconjugated |
|--------------|---|
| Modification | Unmodified |
| lsotype | IgG |
| Clonality | Monoclonal Antibody |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw |
| | cycles. |
| Buffer | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% |
| | BSA |
| Purification | Affinity Purified |

Immunogen

| Gene Name | MAPK14 |
|-------------------|---|
| | MAPK14; CSBP; CSBP1; CSBP2; CSPB1; MXI2; SAPK2A; Mitogen-activated protein kinase |
| Alternative Names | 14; MAP kinase 14; MAPK 14; Cytokine suppressive anti-inflammatory drug-binding |
| | protein; CSAID-binding protein; CSBP; MAP kinase MXI2; MAX-interacting protein |
| Gene ID | 1432 |
| SwissProt ID | Q16539 |

Application

| Dilution Ratio | WB: 1/500-1/1000 IP: 1/20 |
|------------------|--|
| Molecular Weight | Calculated MW: 41 kDa; Observed MW: 41 kDa |




Background

Responds to activation by environmental stress, pro-inflammatory cytokines and lipopolysaccharide (LPS) by phosphorylating a number of transcription factors, such as ELK1 and ATF2 and several downstream kinases, such as MAPKAPK2 and MAPKAPK5. Plays a critical role in the production of some cytokines, for example IL-6.

Research Area

Signal Transduction

Image Data



Western blot analysis of p38 in Hela, A549, MCF-7, U251 lysates using p38 antibody.

Note

For research use only.