

## Summary

Production Name	VAMP8 (8R11) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat

#### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New typepreservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.Avoid freeze / thaw cycle.
Purification	Affinity purification

### Immunogen

Gene Name	VAMP8 {ECO:0000303 PubMed:12130530}
Alternative Names	VAMP8; EDB; Endobrevin; VAMP-8;
Gene ID	8673.0
SwissProt ID	Q9BV40.

# Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	11kDa




#### Background

Proteins in the soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) complex are integral membrane proteins involved in vesicle transport and membrane fusion by pairing of vesicular SNAREs (v-SNAREs) with cognate target SNAREs (t-SNAREs). Vesicle associated membrane protein 8 (VAMP8), also known as endobrevin, is a v-SNARE originally found preferentially localized to early endosomes. SNAREs, soluble N-ethylmaleimide-sensitive factorattachment protein receptors, are essential proteins for fusion of cellular membranes. SNAREs localized on opposing membranes assemble to form a trans-SNARE complex, an extended, parallel four alpha-helical bundle that drives membrane fusion. VAMP8 is a SNARE involved in autophagy through the direct control of autophagosome membrane fusion with the lysososome membrane via its interaction with the STX17-SNAP29 binary t- SNARE complex (PubMed:<a href="http://www.uniprot.org/citations/23217709" target=" blank">23217709</a>, PubMed:<a href="http://www.uniprot.org/citations/25686604" target=" blank">25686604</a>). Also required for dense-granule secretion in platelets (PubMed:<a href="http://www.uniprot.org/citations/12130530" target=" blank">12130530</a>). Plays also a role in regulated enzyme secretion in pancreatic acinar cells (By similarity). Involved in the abscission of the midbody during cell division, which leads to completely separate daughter cells (By similarity). Involved in the homotypic fusion of early and late endosomes (By similarity). Participates also in the activation of type I interferon antiviral response through a TRIM6-dependent mechanism (PubMed: <a href="http://www.uniprot.org/citations/31694946" target=" blank">31694946</a>).

#### **Research Area**

#### Image Data



Western blot analysis of VAMP8 expression in (1) HeLa cell lysate; (2) NIH/3T3 cell lysate; (3) PC-12 cell lysate.

#### Note

For research use only.

